arXiv:1404.7660v1 [math.CA] 30 Apr 2014

SOME NEW GENERAL INTEGRAL INEQUALITIES FOR P-FUNCTIONS

İMDAT İŞCAN▼, ERHAN SET♣, AND M. EMIN ÖZDEMIR■

ABSTRACT. In this paper, we derive new estimates for the remainder term of the midpoint, trapezoid, and Simpson formulae for functions whose derivatives in absolute value at certain power are P-functions. Some applications to special means of real numbers are also given.

1. INTRODUCTION

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a, b \in I$ with $a < b$. The following inequality holds:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a}\int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}. \tag{1.1}$$

This double inequality is well known as Hermite-Hadamard integral inequality for convex functions in the literature .

In [2] Dragomir et al. defined the concept of P-function as the following:

Definition 1. *We say that $f : I \to \mathbb{R}$ is a P-function, or that f belongs to the class $P(I)$, if f is a non-negative function and for all $x, y \in I$, $\alpha \in [0, 1]$, we have*

$$f(\alpha x + (1-\alpha)y) \leq f(x) + f(y).$$

$P(I)$ contain all nonnegative monotone convex and quasi convex functions.

In [2], Dragomir et al., proved following inequalities of Hadamard's type for P-function

Theorem 1. *Let $f \in P(I), a, b \in I$ with $a < b$ and $f \in L[a, b]$. Then the following inequality holds*

$$f\left(\frac{a+b}{2}\right) \leq \frac{2}{b-a}\int_a^b f(x)dx \leq 2\left[f(a)+f(b)\right]. \tag{1.2}$$

The following inequality is well known in the literature as Simpson's inequality .

Date: June 10, 2012.

2000 *Mathematics Subject Classification.* 26A51, 26D15.

Key words and phrases. convex function, P-function, Simpson's inequality, Hermite-Hadamard's inequality.

Let $f:[a,b]\to\mathbb{R}$ be a four times continuously differentiable mapping on (a,b) and $\left\|f^{(4)}\right\|_\infty=\sup_{x\in(a,b)}\left|f^{(4)}(x)\right|<\infty$. Then the following inequality holds:

$$\left|\frac{1}{3}\left[\frac{f(a)+f(b)}{2}+2f\left(\frac{a+b}{2}\right)\right]-\frac{1}{b-a}\int_a^b f(x)dx\right|\leq\frac{1}{2880}\left\|f^{(4)}\right\|_\infty(b-a)^4.$$

In recent years many authors have studied error estimations for Simpson's inequality and Hermite-Hadamard inequalitiy; for refinements, counterparts, generalizations, see ([1]-[10]).

In [3], Iscan obtained a new generalization of some integral inequalities for differentiable convex mapping which are connected Simpson and Hadamard type inequalities by using the following lemma.

Let $f:I\subseteq\mathbb{R}\to\mathbb{R}$ be a differentiable mapping on I° such that $f'\in L[a,b]$, where $a,b\in I$ with $a<b$ and $\alpha,\lambda\in[0,1]$. Then the following equality holds:

$$\begin{aligned}&\lambda\left(\alpha f(a)+(1-\alpha)f(b)\right)+(1-\lambda)f(\alpha a+(1-\alpha)b)-\frac{1}{b-a}\int_a^b f(x)dx\\ =\;&(b-a)\left[\int_0^{1-\alpha}(t-\alpha\lambda)f'(tb+(1-t)a)\,dt\right.\\ &\left.+\int_{1-\alpha}^1(t-1+\lambda(1-\alpha))f'(tb+(1-t)a)\,dt\right].\end{aligned}$$

The aim of this paper is to establish some new general integral inequalities for functions whose derivatives in absolute value at certain power are P-functions. Some applications of these results to special means is to give as well.

2. Main results

Let $f:I\subseteq\mathbb{R}\to\mathbb{R}$ be a differentiable function on I°, the interior of I. Throughout this section we will take

$$\begin{aligned}&I_f(\lambda,\alpha,a,b)\\ =\;&\lambda\left(\alpha f(a)+(1-\alpha)f(b)\right)+(1-\lambda)f(\alpha a+(1-\alpha)b)-\frac{1}{b-a}\int_a^b f(x)dx\end{aligned}$$

where $a,b\in I^\circ$ with $a<b$ and $\alpha,\lambda\in[0,1]$.

Theorem 2. *Let $f:I\subseteq\mathbb{R}\to\mathbb{R}$ be a differentiable mapping on I° such that $f'\in L[a,b]$, where $a,b\in I^\circ$ with $a<b$ and $\alpha,\lambda\in[0,1]$. If $|f'|^q$ is P-function on $[a,b]$, $q\geq1$, then the following inequality holds:*

$$\begin{aligned}(2.1)\qquad&|I_f(\lambda,\alpha,a,b)|\leq(b-a)\left(|f'(b)|^q+|f'(a)|^q\right)^{\frac{1}{q}}\\ &\times\begin{cases}\gamma_2(\alpha,\lambda)+\gamma_2(1-\alpha,\lambda) & \alpha\lambda\leq1-\alpha\leq1-\lambda(1-\alpha)\\ \gamma_2(\alpha,\lambda)+\gamma_1(1-\alpha,\lambda) & \alpha\lambda\leq1-\lambda(1-\alpha)\leq1-\alpha\\ \gamma_1(\alpha,\lambda)+\gamma_2(1-\alpha,\lambda) & 1-\alpha\leq\alpha\lambda\leq1-\lambda(1-\alpha)\end{cases},\end{aligned}$$

where

$$\begin{aligned} \gamma_1(\alpha,\lambda) &= (1-\alpha)\left[\alpha\lambda - \frac{(1-\alpha)}{2}\right], \\ \gamma_2(\alpha,\lambda) &= (\alpha\lambda)^2 - \gamma_1(\alpha,\lambda)\ . \end{aligned} \tag{2.2}$$

Proof. Suppose that $q \geq 1$.Since $|f'|^q$ is P-function on $[a,b]$, from Lemma 1 and using the well known power mean inequality, we have

$$\begin{aligned} &|I_f(\lambda,\alpha,a,b)| \\ &\leq (b-a)\left[\int_0^{1-\alpha} |t-\alpha\lambda|\,|f'(tb+(1-t)a)|\,dt + \int_{1-\alpha}^{1} |t-1+\lambda(1-\alpha)|\,|f'(tb+(1-t)a)|\,dt\right] \\ &\leq (b-a)\left\{\left(\int_0^{1-\alpha} |t-\alpha\lambda|\,dt\right)^{1-\frac{1}{q}}\left(\int_0^{1-\alpha} |t-\alpha\lambda|\,|f'(tb+(1-t)a)|^q\,dt\right)^{\frac{1}{q}}\right. \\ &\quad \left. + \left(\int_{1-\alpha}^{1} |t-1+\lambda(1-\alpha)|\,dt\right)^{1-\frac{1}{q}}\left(\int_{1-\alpha}^{1} |t-1+\lambda(1-\alpha)|\,|f'(tb+(1-t)a)|^q\,dt\right)^{\frac{1}{q}}\right\} \end{aligned}$$

$$\leq (b-a)\left(|f'(b)|^q + |f'(a)|^q\right)^{\frac{1}{q}}\left\{\int_0^{1-\alpha} |t-\alpha\lambda|\,dt + \int_{1-\alpha}^{1} |t-1+\lambda(1-\alpha)|\,dt\right\} \tag{2.3}$$

Additionally, by simple computation

$$\int_0^{1-\alpha} |t-\alpha\lambda|\,dt = \begin{cases} \gamma_2(\alpha,\lambda), & \alpha\lambda \leq 1-\alpha \\ \gamma_1(\alpha,\lambda), & \alpha\lambda \geq 1-\alpha \end{cases}, \tag{2.4}$$

$$\gamma_1(\alpha,\lambda) = (1-\alpha)\left[\alpha\lambda - \frac{(1-\alpha)}{2}\right],\ \gamma_2(\alpha,\lambda) = (\alpha\lambda)^2 - \gamma_1(\alpha,\lambda)\ ,$$

$$\begin{aligned} \int_{1-\alpha}^{1} |t-1+\lambda(1-\alpha)|\,dt &= \int_0^{\alpha} |t-(1-\alpha)\lambda|\,dt \\ &= \begin{cases} \gamma_1(1-\alpha,\lambda), & 1-\lambda(1-\alpha) \leq 1-\alpha \\ \gamma_2(1-\alpha,\lambda), & 1-\lambda(1-\alpha) \geq 1-\alpha \end{cases}, \end{aligned} \tag{2.5}$$

Thus, using (2.4) and (2.5) in (2.3), we obtain the inequality (2.1). This completes the proof. □

Corollary 1. *Under the assumptions of Theorem 2 with $q = 1$, we have*

$$\begin{aligned} &|I_f(\lambda,\alpha,a,b)| \leq (b-a)\left(|f'(b)| + |f'(a)|\right) \\ &\times \begin{cases} \gamma_2(\alpha,\lambda) + \gamma_2(1-\alpha,\lambda) & \alpha\lambda \leq 1-\alpha \leq 1-\lambda(1-\alpha) \\ \gamma_2(\alpha,\lambda) + \gamma_1(1-\alpha,\lambda) & \alpha\lambda \leq 1-\lambda(1-\alpha) \leq 1-\alpha \\ \gamma_1(\alpha,\lambda) + \gamma_2(1-\alpha,\lambda) & 1-\alpha \leq \alpha\lambda \leq 1-\lambda(1-\alpha) \end{cases}, \end{aligned}$$

Corollary 2. *In Theorem 2 , if we take* $\alpha = \frac{1}{2}$ *and* $\lambda = \frac{1}{3}$, *then we have the following Simpson type inequality*

$$\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{1}{b-a}\int_a^b f(x)dx\right| \leq \frac{5\,(b-a)}{36}\left(|f'(b)|^q+|f'(a)|^q\right)^{\frac{1}{q}}$$

Corollary 3. *In Theorem 2 , if we take* $\alpha = \frac{1}{2}$ *and* $\lambda = 0$, *then we have following midpoint inequality*

$$\left|f\left(\frac{a+b}{2}\right)-\frac{1}{b-a}\int_a^b f(x)dx\right| \leq \frac{b-a}{4}\left(|f'(b)|^q+|f'(a)|^q\right)^{\frac{1}{q}}$$

Corollary 4. *In Theorem 2 , if we take* $\alpha = \frac{1}{2}$ *, and* $\lambda = 1$, *then we get the following trapezoid inequality which is identical to the inequality in* [1, Theorem 2.3].

$$\left|\frac{f\,(a)+f\,(b)}{2}-\frac{1}{b-a}\int_a^b f(x)dx\right| \leq \frac{b-a}{4}\left(|f'(b)|^q+|f'(a)|^q\right)^{\frac{1}{q}}$$

Using Lemma 1 we shall give another result for convex functions as follows.

Theorem 3. *Let* $f : I \subseteq \mathbb{R} \to \mathbb{R}$ *be a differentiable mapping on* I° *such that* $f' \in L[a,b]$, *where* $a, b \in I^\circ$ *with* $a < b$ *and* $\alpha, \lambda \in [0,1]$. *If* $|f'|^q$ *is P-function on* $[a,b]$, $q > 1$, *then the following inequality holds:*

$$|I_f\,(\lambda,\alpha,a,b)| \leq (b-a)\left(\frac{1}{p+1}\right)^{\frac{1}{p}} \tag{2.6}$$

$$\times \begin{cases} \left[\varepsilon_1^{1/p}(\alpha,\lambda,p)c_f^{1/q}(\alpha,q)+\varepsilon_1^{1/p}(1-\alpha,\lambda,p)k_f^{1/q}(\alpha,q)\right], & \alpha\lambda \leq 1-\alpha \leq 1-\lambda\,(1-\alpha) \\ \left[\varepsilon_1^{1/p}(\alpha,\lambda,p)c_f^{1/q}(\alpha,q)+\varepsilon_2^{1/p}(1-\alpha,\lambda,p)k_f^{1/q}(\alpha,q)\right], & \alpha\lambda \leq 1-\lambda\,(1-\alpha) \leq 1-\alpha \\ \left[\varepsilon_2^{1/p}(\alpha,\lambda,p)c_f^{1/q}(\alpha,q)+\varepsilon_1^{1/p}(1-\alpha,\lambda,p)k_f^{1/q}(\alpha,q)\right], & 1-\alpha \leq \alpha\lambda \leq 1-\lambda\,(1-\alpha) \end{cases},$$

where

$$\begin{aligned} c_f(\alpha,q) &= (1-\alpha)\left[|f'\left((1-\alpha)\,b+\alpha a\right)|^q+|f'\,(a)|^q\right], \\ k_f(\alpha,q) &= \alpha\left[|f'\left((1-\alpha)\,b+\alpha a\right)|^q+|f'\,(b)|^q\right], \end{aligned} \tag{2.7}$$

$\varepsilon_1(\alpha,\lambda,p) = (\alpha\lambda)^{p+1}+(1-\alpha-\alpha\lambda)^{p+1}$, $\varepsilon_2(\alpha,\lambda,p) = (\alpha\lambda)^{p+1}-(\alpha\lambda-1+\alpha)^{p+1}$, *and* $\frac{1}{p}+\frac{1}{q} = 1$.

Proof. Since $|f'|^q$ is P-function on $[a,b]$, from Lemma 1 and by Hölder's integral inequality, we have

$$\begin{aligned} & |I_f\,(\lambda,\alpha,a,b)| \\ \leq\ & (b-a)\left[\int_0^{1-\alpha}|t-\alpha\lambda|\,|f'\,(tb+(1-t)a)|\,dt+\int_{1-\alpha}^1|t-1+\lambda\,(1-\alpha)|\,|f'\,(tb+(1-t)a)|\,dt\right] \end{aligned}$$

$$\leq (b-a)\left\{\left(\int_0^{1-\alpha}|t-\alpha\lambda|^p\,dt\right)^{\frac{1}{p}}\left(\int_0^{1-\alpha}|f'(tb+(1-t)a)|^q\,dt\right)^{\frac{1}{q}}\right.$$
$$\left.+\left(\int_{1-\alpha}^{1}|t-1+\lambda(1-\alpha)|^p\,dt\right)^{\frac{1}{p}}\left(\int_{1-\alpha}^{1}|f'(tb+(1-t)a)|^q\,dt\right)^{\frac{1}{q}}\right\}. \tag{2.8}$$

By the inequality (1.2), we get

$$\int_0^{1-\alpha}|f'(tb+(1-t)a)|^q\,dt = (1-\alpha)\left[\frac{1}{(1-\alpha)(b-a)}\int_a^{(1-\alpha)b+\alpha a}|f'(x)|^q\,dx\right]$$
$$\leq (1-\alpha)\left[|f'((1-\alpha)b+\alpha a)|^q+|f'(a)|^q\right]. \tag{2.9}$$

The inequality (2.9) also holds for $\alpha=1$. Similarly, for $\alpha\in(0,1]$ by the inequality (1.2), we have

$$\int_{1-\alpha}^{1}|f'(tb+(1-t)a)|^q\,dt = \alpha\left[\frac{1}{\alpha(b-a)}\int_{(1-\alpha)b+\alpha a}^{b}|f'(x)|^q\,dx\right]$$
$$\leq \alpha\left[|f'((1-\alpha)b+\alpha a)|^q+|f'(b)|^q\right]. \tag{2.10}$$

The inequality (2.10) also holds for $\alpha=0$. By simple computation

$$\int_0^{1-\alpha}|t-\alpha\lambda|^p\,dt=\begin{cases}\frac{(\alpha\lambda)^{p+1}+(1-\alpha-\alpha\lambda)^{p+1}}{p+1}, & \alpha\lambda\leq 1-\alpha\\ \frac{(\alpha\lambda)^{p+1}-(\alpha\lambda-1+\alpha)^{p+1}}{p+1}, & \alpha\lambda\geq 1-\alpha\end{cases}, \tag{2.11}$$

and

$$\int_{1-\alpha}^{1}|t-1+\lambda(1-\alpha)|^p\,dt=\begin{cases}\frac{[\lambda(1-\alpha)]^{p+1}+[\alpha-\lambda(1-\alpha)]^{p+1}}{p+1}, & 1-\alpha\leq 1-\lambda(1-\alpha)\\ \frac{[\lambda(1-\alpha)]^{p+1}-[\lambda(1-\alpha)-\alpha]^{p+1}}{p+1}, & 1-\alpha\geq 1-\lambda(1-\alpha)\end{cases}, \tag{2.12}$$

thus, using (2.9)-(2.12) in (2.8), we obtain the inequality (2.6). This completes the proof. □

Corollary 5. *In Theorem 3, if we take $\alpha=\frac{1}{2}$ and $\lambda=\frac{1}{3}$, then we have the following Simpson type inequality*

$$\left|\frac{1}{6}\left[f(a)+4f\left(\frac{a+b}{2}\right)+f(b)\right]-\frac{1}{b-a}\int_a^b f(x)dx\right|\leq\frac{b-a}{12}\left(\frac{1+2^{p+1}}{3(p+1)}\right)^{\frac{1}{p}}$$
$$\times\left\{\left(\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(a)|^q\right)^{\frac{1}{q}}+\left(\left|f'\left(\frac{a+b}{2}\right)\right|^q+|f'(b)|^q\right)^{\frac{1}{q}}\right\}.$$

Corollary 6. *In Theorem 3, if we take $\alpha = \frac{1}{2}$ and $\lambda = 0$, then we have the following midpoint inequality*

$$\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a}\int_a^b f(x)dx \right| \leq \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}} \times \left\{ \left(\left| f'\left(\frac{a+b}{2}\right)\right|^q + |f'(a)|^q \right)^{\frac{1}{q}} + \left(\left| f'\left(\frac{a+b}{2}\right)\right|^q + |f'(b)|^q \right)^{\frac{1}{q}} \right\}.$$

We note that by inequality

$$\left| f'\left(\frac{a+b}{2}\right)\right|^q \leq |f'(a)|^q + |f'(b)|^q$$

we have

$$\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a}\int_a^b f(x)dx \right| \leq \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}} \left\{ \left(|f'(b)|^q + 2|f'(a)|^q\right)^{\frac{1}{q}} + \left(|f'(a)|^q + 2|f'(b)|^q\right)^{\frac{1}{q}} \right\}.$$

Corollary 7. *In Theorem 3, if we take $\alpha = \frac{1}{2}$ and $\lambda = 1$, then we have the following trapezoid inequality*

$$\left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)dx \right| \leq \frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}} \times \left\{ \left(\left| f'\left(\frac{a+b}{2}\right)\right|^q + |f'(a)|^q \right)^{\frac{1}{q}} + \left(\left| f'\left(\frac{a+b}{2}\right)\right|^q + |f'(b)|^q \right)^{\frac{1}{q}} \right\}.$$

3. Some applications for special means

We now recall the following well-known concepts. For arbitrary real numbers a, b, $a \neq b$, we define

(1) The unweighted arithmetic mean

$$A(a,b) := \frac{a+b}{2},\ a,b \in \mathbb{R}.$$

(2) Then $n-$Logarithmic mean

$$L_n(a,b) := \left(\frac{b^{n+1}-a^{n+1}}{(n+1)(b-a)}\right)^{\frac{1}{n}},\ n \in \mathbb{N},\ n \geq 1,\ a,b \in \mathbb{R},\ a < b.$$

Now we give some applications of the new results derived in section 2 to special means of real numbers.

Proposition 1. *Let $a, b \in \mathbb{R}$ with $a < b$ and $n \in \mathbb{N}, n \geq 2$. Then*

$$\left| \frac{1}{3}A(a^n, b^n) + \frac{2}{3}A^n(a,b) - L_n^n(a,b) \right| \leq \frac{5n(b-a)}{36}\left(|b|^{(n-1)q} + |a|^{(n-1)q}\right)^{\frac{1}{q}}$$

Proof. The assertion follows from Corollary 2 applied to the function $f(x) = x^n$, $x \in \mathbb{R}$, because $|f'|^q$ is a P-function. □

Proposition 2. *Let $a,b\in\mathbb{R}$ with $a<b$ and $n\in\mathbb{N},n\geq 2$. Then*

$$|A^n(a,b)-L_n^n(a,b)|\leq\frac{n\,(b-a)}{4}\left(|b|^{(n-1)q}+|a|^{(n-1)q}\right)^{\frac{1}{q}}$$

and

$$|A(a^n,b^n)-L_n^n(a,b)|\leq\frac{n\,(b-a)}{4}\left(|b|^{(n-1)q}+|a|^{(n-1)q}\right)^{\frac{1}{q}}$$

Proof. The assertion follows from Corollary 3 and Corollary 4 applied to the function $f(x)=x^n$, $x\in\mathbb{R}$, because $|f'|^q$ is a P-function. □

Proposition 3. *Let $a,b\in\mathbb{R}$ with $a<b$ and $n\in\mathbb{N},n\geq 2$. Then*

$$\left|\frac{1}{3}A(a^n,b^n)+\frac{2}{3}A^n(a,b)-L_n^n(a,b)\right|\leq\frac{n\,(b-a)}{12}\left(\frac{1+2^{p+1}}{3\,(p+1)}\right)^{\frac{1}{p}}$$
$$\times\left\{\left(|A(a,b)|^{(n-1)q}+|a|^{(n-1)q}\right)^{\frac{1}{q}}+\left(|A(a,b)|^{(n-1)q}+|b|^{(n-1)q}\right)^{\frac{1}{q}}\right\}.$$

Proof. The assertion follows from Corollary 5 applied to the function $f(x)=x^n$, $x\in\mathbb{R}$, because $|f'|^q$ is a P-function. □

Proposition 4. *Let $a,b\in\mathbb{R}$ with $a<b$ and $n\in\mathbb{N},n\geq 2$. Then*

$$|A^n(a,b)-L_n^n(a,b)|\leq\frac{n\,(b-a)}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}$$
$$\times\left\{\left(|A(a,b)|^{(n-1)q}+|a|^{(n-1)q}\right)^{\frac{1}{q}}+\left(|A(a,b)|^{(n-1)q}+|b|^{(n-1)q}\right)^{\frac{1}{q}}\right\}.$$

and

$$|A(a^n,b^n)-L_n^n(a,b)|\leq\frac{b-a}{4}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}$$
$$\times\left\{\left(|A(a,b)|^{(n-1)q}+|a|^{(n-1)q}\right)^{\frac{1}{q}}+\left(|A(a,b)|^{(n-1)q}+|b|^{(n-1)q}\right)^{\frac{1}{q}}\right\}.$$

Proof. The assertion follows from Corollary 6 and Corollary 7 applied to the function $f(x)=x^n$, $x\in\mathbb{R}$, because $|f'|^q$ is a P-function. □

References

[1] A. Barani and S. Barani, Hermite-Hadamard type inequalities for functions when a power of the absolute value of the first derivative is P-convex, Bull. Aust. Math. Soc., 86 (1) (2012), 126-134.

[2] S.S. Dragomir, J. Pečarić, L.E. Persson, Some inequalities of Hadamard type, Soochow J. Math. **21** (1995), 335-341.

[3] İ. İşcan, A new generalization of some integral inequalities and their applications, International Journal of Engineering and Applied sciences, **3**(3) (2013), 17-27.

[4] İ. İşcan, Some new general integral inequalities for h-convex and h-concave functions, Advances in Pure and Applied Mathematics, 2014. DOI: 10.1515/apam-2013-0029.

[5] U.S. Kırmacı, Inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula, Appl. Math. Comp., 147 (2004), 137-146.

[6] U.S. Kırmacı, M.E. Özdemir, On some inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula, Appl. Math. Comp., 153 (2004), 361-368.

[7] M. E. Özdemir and Ç. Yıldız, New inequalities for Hermite-Hadamard and Simpson type with applications, Tamkang journal of Mathematics, 44 (2) (2013), 209-216.

[8] M.Z. Sarıkaya, E. Set, M. E. Özdemir, On new inequalities of Simpson's type for s-convex functions, Comp. Math. Appl., 60 (2010), 2191-2199.
[9] E. Set, M. E. Özdemir, M.Z. Sarıkaya, On new inequalities of Simpson's type for quasi-convex functions with applications, Tamkang J. Math., 43 (3) (2012), 357-364.
[10] M. Tunc, Ç. Yıldız, A. Ekinci, On some inequalitiesof Simpson's type via h-convex functions, Hacettepe Journal of Mathematics and Statistics, 42 (4) (2013), 309-317.

[▼]Department of Mathematics, Faculty of Arts and Sciences, Giresun University, 28100, Giresun, Turkey.
E-mail address: imdat.iscan@giresun.edu.tr, imdati@yahoo.com

[♣]Department of Mathematics, Faculty of Arts and Sciences, Ordu University, 52200, Ordu, Turkey
E-mail address: erhanset@yahoo.com

[■]Atatürk University, K.K. Education Faculty, Department of Mathematics, 25240, Campus, Erzurum, Turkey
E-mail address: emos@atauni.edu.tr